# Psychological and Social Factors Affecting Sexual Desire in Adult Relationships

Posted on July 3, 2018

View this essay online:
https://academic-master.com/psychological-and-social-factors-affecting-sexual-desire-in-adult-relationships/

## Introduction

Sexual desire changes across adulthood and is influenced by physical health, medication, mood, sleep, stress, relationship quality, substance use, privacy, culture, and personal meaning. A clinically important change is usually one that is persistent, distressing, or linked with another medical or psychological problem rather than simply different from a partner's preferred frequency. Two cases illustrate this complexity. Veronica, a fifty-seven-year-old man, reports reduced desire in the context of diabetes, hypertension, stress, depression, and marijuana use. Jorge, a nineteen-year-old heterosexual man, notices lower libido and sexual-performance difficulty after beginning a selective serotonin reuptake inhibitor for depression. Neither case should be framed as a failure to satisfy another person or to meet a cultural standard of masculinity. Sexual activity requires voluntary consent, and treatment should focus on the individual's wellbeing, relationship communication, and informed choice. Assessment is strongest when biological, psychological, and social explanations are considered together rather than when one symptom is attributed automatically to age, medication, orientation, or relationship status.

## Veronica's Medical and Psychological Picture

Veronica's diabetes and hypertension deserve careful attention because both can affect vascular function, nerve function, energy, mood, and erectile performance. Low desire and erectile dysfunction are related but distinct: reduced libido concerns interest in sexual activity, while erectile dysfunction concerns difficulty obtaining or maintaining an erection adequate for desired sexual activity. A clinician should therefore ask about morning erections, orgasm, ejaculation, pain, masturbation, relationship-specific patterns, and the timing of symptoms. Medication review is also essential because some antidepressants, antihypertensives, opioids, and sedatives can affect desire or performance. Depression, sleep loss, chronic stress, and body-image concerns may reduce pleasure independently of medical illness. Laboratory testing may be appropriate when the history suggests low testosterone, thyroid disease, elevated prolactin, or another endocrine problem, but symptoms alone do not establish a hormone deficiency. The goal is to identify modifiable contributors while avoiding unsupported "testosterone boosters" or unverified sexual-enhancement products.

# Relationship Ethics and Sexual Communication

Relationship tension can intensify low desire when one partner experiences reduced interest as rejection and the other experiences requests for sex as pressure or evaluation. Marriage does not create an obligation to provide sex, and no clinician should recruit one partner to pressure the other into intercourse. A more useful approach is to identify what each person means by intimacy, affection, initiation, and satisfaction. Couples may benefit from discussing timing, privacy, nonsexual touch, expectations, and activities that feel comfortable without assuming that every affectionate interaction must lead to intercourse. Therapy focused on sexual health can help reduce performance anxiety and the pursue-withdraw cycle that develops when one partner repeatedly initiates and the other increasingly avoids intimacy. Pornography or sexual media should not be prescribed as a universal remedy; some adults find it acceptable, while others find it inconsistent with personal values or relationship boundaries. Consent, mutual preference, and communication matter more than any single technique.

# Jorge, Depression, and SSRI Sexual Side Effects

Jorge's case requires separating depression itself from the effects of antidepressant treatment. Depression can reduce energy, pleasure, self-confidence, and sexual interest, but SSRIs can also reduce libido, delay orgasm or ejaculation, and contribute to erectile difficulty. MedlinePlus lists sexual problems, including loss of desire or ability to have sex, among common antidepressant side effects and advises patients not to change the dose or stop treatment on their own. The timing of Jorge's symptoms therefore matters: clinicians should ask about sexual function before medication, the onset after treatment began or the dose changed, improvement in depressive symptoms, and the specific type of sexual difficulty. Depending on severity and treatment response, the prescriber may consider watchful waiting, dose adjustment, timing changes, switching medication, or another strategy. Jorge's heterosexuality and consensual casual relationships are not disorders. Peer pressure, however, can worsen anxiety by treating sexual activity as proof of masculinity. Treatment should reflect Jorge's own goals rather than his friends' expectations.

# Substances, Lifestyle, and Preventive Sexual Health

Cannabis, alcohol, tobacco, sleep, exercise, and chronic stress can all influence sexual function, but effects differ among individuals. Veronica's marijuana use should be discussed without assuming either benefit or harm. Some people report relaxation or altered sensation, while others experience anxiety, reduced motivation, or impaired performance. Alcohol may reduce inhibition while also impairing arousal, and tobacco damages vascular health. Regular physical activity, adequate sleep, improved diabetes management, and treatment of depression can support general health even when

a separate sexual-function intervention is still needed. Preventive care is equally important for Jorge if he has multiple partners. Condoms or other barrier methods, vaccination, STI testing based on exposure, contraception discussions with partners, and HIV prevention when appropriate should be addressed without moral judgment. Consent must remain clear when alcohol or drugs are involved. Sexual-health counseling works best when patients can disclose behavior honestly without fear that clinicians will equate orientation, partner number, or reduced desire with pathology.

# A Biopsychosocial Treatment Strategy

A practical plan begins by defining the exact concern and then matching treatment to its likely contributors. For Veronica, priorities may include improving blood-pressure and diabetes control, reviewing medication, screening for sleep problems and depression, evaluating possible endocrine causes when indicated, and addressing relationship stress. If erectile dysfunction is present, evidence-based treatment can be discussed separately from desire, since erection-enhancing medications do not necessarily increase libido and may be unsafe with certain cardiovascular drugs such as nitrates. For Jorge, the prescriber should weigh antidepressant benefit against sexual side effects while monitoring depression and suicide risk. Both men may benefit from counseling that reduces shame and performance pressure. Treatment success should not be measured only by intercourse frequency. Better mood, restored desire, improved communication, satisfying nonpenetrative intimacy, safer behavior, fewer medication side effects, and reduced distress may all represent meaningful improvement. Because sexual function changes over time, goals and treatment should be reviewed rather than treated as one permanent prescription.

# Conclusion

Sexual desire is shaped by interacting medical, psychological, relational, and social factors rather than by one fixed biological drive. Veronica’s diabetes, hypertension, depression, stress, medicines, and marijuana use all deserve structured assessment, while relationship counseling should protect consent rather than frame sex as a marital duty. Jorge’s symptoms are plausibly related to both depression and SSRI treatment, and current evidence supports taking antidepressant sexual side effects seriously rather than dismissing them. Neither man should be judged according to another person’s preferred frequency of sex, cultural expectations of masculinity, or moral assumptions about consensual adult behavior. Evidence-based care identifies the specific problem, reviews medical and medication contributors, addresses mood and sleep, supports communication, and includes preventive sexual-health information when relevant. The most appropriate goal is not performance for its own sake but improved wellbeing, autonomy, safety, and satisfaction as defined by the patient and, where relational goals are involved, developed collaboratively with a consenting partner.

# References

American Psychiatric Association. (2022). *Diagnostic and Statistical Manual of Mental Disorders* (5th ed., text rev.).

McCabe, M. P., et al. (2016). Risk factors for sexual dysfunction among women and men. *Journal of Sexual Medicine, 13*(2), 153–167.

MedlinePlus. (2026). *Antidepressants*.

National Institute of Diabetes and Digestive and Kidney Diseases. (2024). *Symptoms, causes, diagnosis, and treatment of erectile dysfunction*.