

Types of Erectile Dysfunction and Associated Medications and Therapies

Posted on May 8, 2018

Erectile dysfunction is the failure to get or keep an erection sufficiently firm to have sex. (National Institute of Diabetes and Digestive and Kidney Diseases [NIDDK], n.d.-a) It is also referred to as impotence in medical terminology. Occasional ED is also not uncommon. Many men experience occasional erectile dysfunction during times of stress. Frequent erectile dysfunction can be a sign of health problems that can be overcome by treatment. It is also a signal of relationship or emotional difficulty that may require to be reported to a doctor. Not all male sexual problems are caused by erectile dysfunction. Other types of male sexual dysfunction include the following: premature ejaculation, lack of interest in sex, and delayed or absent ejaculation. Frequent erectile dysfunction report trouble getting an erection, reduced interest in sex, and Difficulty maintaining an erection during sexual activities Other sexual disorders related to ED include delayed ejaculation, premature ejaculation, and an orgasm, which is the failure to achieve orgasm after full stimulation.

Types of erectile dysfunction:

Erectile dysfunction isn't the same as another. Specialists make qualifications between various kinds of weaknesses:

Primacy dysfunction:

In this condition, the man has never had an erection and will never accomplish one suddenly. This is generally caused by a physical variation from the norm in the penis, for example, a caught nerve. Essential feebleness can now and then be treated with surgery.

Secondary dysfunction:

Men who experience the ill effects of this have had full erections previously; however, have experienced erectile brokenness sooner or later.

Situational impotence:

This implies the man figures out how to have a full erection in specific circumstances. He can accomplish an erection amid masturbation, for instance, but not amid sex with an accomplice.

Incomplete impotence:

In this erectile dysfunction, the penis turns out to be hardened, however soon thereafter winds up plainly floppy. It is likewise conceivable that the penis progresses toward becoming semi-unbending

Total impotence:

This term demonstrates that there are no erections by any stretch of the imagination. Not by any means early morning erections.

How erectile dysfunction affects:

Erectile dysfunction psychologically affects most men, notwithstanding when the reason for the issue is absolutely a physical one. Erectile dysfunction can likewise cause relationship issues.

Common feelings of men with Erectile Dysfunction are mentioned:

- Low self-esteem
- Feeling 'loss of manhood'
- Sad about his loss
- Scared of approaching a sexual partner
- Feelings of insecurity
- Low self-esteem
- Frightened about being intimate with a partner
- Angry and aggressive
- Feeling ashamed
- Worry about not being competent to pleasure their partner
- Feeling all-time guilty about their dysfunction

Normal emotions accomplices of men with Erectile Dysfunction encounter are:

- Blameworthy about not having the capacity to influence their accomplice's penis to erect
- Powerless, not certain where to go for help
- Rejected/Unloved, not certain if their accomplice is still into them
- Feeling ugly

- Befuddled/Frustrated
- Terrified/stressed that he is engaging in extramarital relations
- Stressed over the well-being and prosperity of their friends and family
- Unreliable about his affection
- Dismal for him, or notwithstanding for themselves, about not having an erect penis in the room any longer
- A few accomplices, ladies specifically, are alleviated their man's penis isn't all they bring into the room any longer

Treatment of erectile dysfunction:

A large number of men after the age of 50 experience the ill effects of ED or erectile brokenness. A large number of these men have endorsed medications, for example, Viagra, Levitra, or Cialis which are regularly effective. Be that as it may, men can adopt a non-restorative strategy to fathoming erectile dysfunction. By taking five action steps any man with ED can go out on a limb and even treat concerns of ED.

Five action steps can help men with ED. (NIDDK, n.d.-b) These are appropriate weight administration, sufficient rest, an organized exercise program, restricted liquor consumption and a legitimate, adjusted eating regimen.

Here, their men need to roll out improvements in their ways of life. Sexual brokenness in men is regularly caused by something that is likewise perilous, for example, cardiovascular issues. Men who make these ways of life move steps are frequently ready to go off of drugs including expanding their sexual execution.

- Adequate sleep
- Weight management
- Balanced diet
- Regular sleep
- And limited alcohol

For some men, oral solutions are exceptionally protected, as well as very compelling. Pills, for example, Viagra, Levitra, and Cialis are viable for around 70% of men. Be that as it may, for those with other well-being factors, for example, heart ailment, or diabetes, the oral solution may not be a choice, or they may turn out to be less viable after some time. For these men, there are various other treatment alternatives accessible, including:

Vacuum pump therapy:

This requires the utilization of a little pump to upgrade blood flow and make an erection mechanically.

These gadgets are absolutely noninvasive, yet they can be to some degree ungainly to utilize and not helpful for suddenness.

Trans urethral suppositories:

This treatment involves small pellets of medicine embedded into the urethra opening – a negligibly obtrusive treatment, yet at the same time to some degree uncomfortable. This likewise requires some propel arrangement.

Injection therapy:

while the idea is exasperating to most men toward the beginning, includes the utilization of a fine needle to infuse medicine to deliver an erection. Due to both the [mental and physical](#) inconvenience, numerous men don't observe this to be a long-haul arrangement regardless of large amounts of viability. Around half of the men who attempt infusions, don't proceed with their utilization.

Prosthetic inserts:

These are surgical instruments disguised inside the body that reestablish erectile capacity to as close to normal as could be expected under the circumstances and still take into account immediacy. While the idea is just the same old thing new, the gadgets and implantation methods have advanced significantly to make the system speedier and less demanding on the patient. While inserts convey similar dangers related to any surgery, embed methodology, for the most part, has a great degree of high achievement rates and gives a viable answer for some men. Inserts may not be for everybody, but rather, they are a feasible alternative to reestablish erectile capacity for most men beyond 50 years old, they convey a high fulfilment rate among patients and accomplices.

An expected 30 million men have erectile dysfunction (NIDDK, n.d.-a), and seven out of 10 cases are caused by a conceivably lethal condition like atherosclerosis, kidney sickness, vascular disease, neurological problems, or diabetes. Specific solutions can likewise cause erectile dysfunction, surgical methodology on the prostate organ and colon, and mental health issues. Here's a glance at the confirmation for and against six of the most prominent regular [cures or herbal](#) remedies:

- Arginine,
- Ginseng,
- Pomegranate juice,
- Yohimbe,
- Horny goat weed,
- Ginkgo Biloba.

Doctors endorse prescriptions to help deal with the side effects of Erectile dysfunction. You may need

to attempt a few meds before discovering one that works. These solutions can have symptoms. In case you're encountering obnoxious symptoms, converse with your specialist. They might have the capacity to prescribe an alternate prescription.

The following medications stimulate blood flow to your penis to help treat ED:

Alprostadil, Avanafil, Sildenafil, Cialis, Androderm, Levitra. (NIDDK, n.d.-b)

If erectile dysfunction is affecting your relationship, you may also think of speaking with a relationship counsellor. Relationship counselling will help reconnect emotionally with a partner, which may also help to recover erectile dysfunction. (NIDDK, n.d.-b)

References:

Araujo, A.B., Durante, R., Feldman, H.A. The relationship between depressive symptoms and male erectile dysfunction: cross-sectional results from the Massachusetts Male Aging Study. *Psychosom Med.* 1998;60:458-465.

Derogatis, L.R., Meyer, J.K., King, K.M. Psychopathology in individuals with sexual dysfunction. *Am J Psychiatry.* 1981;138:757-763.

Schreiner-Engel, P., Schiavi, R.C. Lifetime psychopathology in individuals with low sexual desire. *J Nerv Ment Dis.* 1986;174:646-651.

National Institute of Diabetes and Digestive and Kidney Diseases. (n.d.-a). *Definition and facts for erectile dysfunction*. National Institutes of Health.
<https://www.niddk.nih.gov/health-information/urologic-diseases/erectile-dysfunction/definition-facts>

National Institute of Diabetes and Digestive and Kidney Diseases. (n.d.-b). *Treatment for erectile dysfunction*. National Institutes of Health.
<https://www.niddk.nih.gov/health-information/urologic-diseases/erectile-dysfunction/treatment>

National Institute of Diabetes and Digestive and Kidney Diseases. (n.d.-c). *Symptoms and causes of erectile dysfunction*. National Institutes of Health.
<https://www.niddk.nih.gov/health-information/urologic-diseases/erectile-dysfunction/symptoms-causes>